

### Relationship of HIV Disease Cases, HIV Cases, and AIDS Cases




From the time a person is infected with human immunodeficiency virus (HIV) until death, he/she has **HIV Disease**. All persons with HIV Disease can be subclassified as either an **AIDS case** (if they are in the later stages of the disease process and have met the case definition for AIDS) or an **HIV case** (if they are in the earlier stages of the disease process and have not met the AIDS case definition).



<sup>\*</sup> HIV Case vs. AIDS Case


These cases were initially reported as HIV cases, and have subsequently remained HIV cases (i.e., they have not met the case definition for AIDS)



<sup>\*</sup> Based on death certificate data.

<sup>\*\*</sup>Cases are indicated by year of their initial report to the Missouri Department of Health and Senior Services(i.e., by the year in which the first report of the person, whether as an HIV case or an AIDS case, was received by the department.)

These cases were either: 1) initially reported as HIV cases and then later reclassified as AIDS cases because they had subsequently come to meet the AIDS case definition: or 2) initially reported as an AIDS case.




## Reported HIV and AIDS Cases by Gender, Race/Ethnicity, and Age at Diagnosis, Missouri, 1982–2001


	HIV Cases*			AIDS C	ases**		HIV/AIDS Cases		
Repo	Reported 2001		ulative*	Repor	rted 2001	Cum	Cumulative Cum		lative
Case	s %	Cases	%	Cases	%	Cases	%	Cases	%
Gender									
Male		3,756	(82.9%)	301	(81.4%)	8,225	(90.2%).	11,981	(87.8%)
Female96	(22.7%)	776	(17.1%)	69	(18.6%)	894	(9.8%).	1,670	(12.2%)
Race/Ethnicity									
White	(47.6%)	2,445	(53.9%)	160	(43.2%)	5,914	(64.9%).	8,359	(61.2%)
Black	(47.9%)	1,935	(42.7%)	197	(53.2%)	2,966	(32.5%).	4,901	(35.9%)
Hispanic10	(2.4%)	104	(2.3%)		(2.2%)	182	(2.0%).	286	(2.1%)
Asian/PacificIslander1	(0.2%)	14	(0.3%)			25	, ,	39	(0.3%)
American Indian0		13	(0.3%)		,	32	, ,	45	(0.3%)
Unknown8	(1.9%)	21	(0.5%)	0	(0.0%)	0	(0.0%).	21	(0.2%)
Race/Ethnicity and Gender									
White Male 173	(41.0%)	2,152	(47.5%)	142	(38.4%)	5,535	(60.7%).	7,687	(56.3%)
BlackMale 139	(32.9%)	1,475	(32.5%)	148	(40.0%)	2,470		3,945	(28.9%)
Hispanic Male7	(1.7%)	90	(2.0%)			169	( , .	259	(1.9%)
Asian/Pacific Islander Male1	(0.2%)	10	(0.2%)			22		32	(0.2%)
American Indian Male0	(0.0%)	12	(0.3%)		,	29		41	(0.3%)
Unknown Male6	(1.4%)	17	(0.4%)	0	, ,	0	(0.0%).	17	(0.1%)
WhiteFemale28	(6.6%)	293	(6.5%)			379		672	(4.9%)
BlackFemale63	(14.9%)	460	(10.2%)		( ,	496		956	(7.0%)
Hispanic Female3		14	(0.3%)			13	(	27	(0.2%)
Asian/Pacific Islander Female0	(/	4	(0.1%)			3		7	(0.1%)
American Indian Female0	(	1	(0.0%)		, ,	3	, ,	4	(0.0%)
Unknown Female2	(0.5%)	4	(0.1%)	0	(0.0%)	0	(0.0%).	4	(0.0%)
Age at Diagnosis‡									
<134	(0.9%)	45	(1.0%)	1	(0.3%)	57	(0.6%).	102	(0.7%)
13-1918	(4.3%)	210	(4.6%)	4	(1.1%)	96	(1.1%).	306	(2.2%)
20-2456	(13.3%)	704	(15.6%)	15	(4.1%)	542	(5.9%).	1,246	(9.1%)
25-2965	(15.4%)	995	(22.0%)	40	(10.8%)	1,486		2,481	(18.2%)
30-3458	(13.7%)	989	(21.8%)		,	2,265		3,254	(23.8%)
35-3991	(21.6%)	731	(16.1%)		( ,	1,892		2,623	(19.2%)
40-44	(13.7%)	437	(9.6%)			1,250		1,687	(12.4%)
45-49	(9.0%)	226	(5.0%)		,	727	` /	953	(7.0%)
50+34	(8.1%)	195	(4.3%)	47	(12.7%)	804	(8.8%).	999	(7.3%)
Missouri Total422	(100.0%)	4,532	(100.0%)	370	(100.0%)	9,119	(100.0%)	13,651	(100.0%)


<sup>\*</sup>HIV Cases-Persons with HIV infection who have not developed one of the specific diseases or conditions which would cause them to meet the case definition for AIDS.
\*\*AIDS Cases-Persons with HIV infection who have developed one or more of the specific diseases or conditions which cause them to meet the AIDS case definition.

<sup>&</sup>lt;sup>‡</sup>For HIV Cases, Age at Diagnosis is the age at which the individual was first diagnosed with HIV infection.

For AIDS Cases, Age at Diagnosis is the age at which the individual was first diagnosed with AIDS.







### Reported HIV and AIDS Cases by Exposure Category, Missouri, 1982–2001


	HIV Cases*				AIDS Cases**				<b>HIV/AIDS Cases</b>	
Rep	orted 2001	Cum	<u>Cumulative</u> Re		eported 2001 Cu		nulative	Cumu	Cumulative	
Cas	es %	Cases	%	Cases	%	Cases	%	Cases	%	
Exposure Category <sup>¶</sup>										
MSM 166	(39.3%)	2,621	(57.8%)	194	(52.4%)	6,335	(69.5%)	8,956	(65.6%)	
MSM/IDU10	(2.4%)	265	(5.8%)	17	(4.6%)	794	(8.7%)	1,059	(7.8%)	
IDU18	(4.3%)	392	(8.6%)	37	(10.0%)	680	(7.5%)	1,072	(7.9%)	
Heterosexual Contact78	(18.5%)	748	(16.5%)	68	(18.4%)	792	(8.7%)	1,540	(11.3%)	
Adult Hemophiliac	(0.2%)	30	(0.7%)	1	(0.3%)	145	(1.6%)	175	(1.3%)	
Adult Transfusion	2 (0.5%)	15	(0.3%)	3	(0.8%)	101	(1.1%)	116	(0.8%)	
Other/Unknown Adult 143	(33.9%)	416	(9.2%)	48	(13.0%)	204	(2.2%)	620	(4.5%)	
Perinatal Transmission	(0.7%)	37	(0.8%)	1	(0.3%)	46	(0.5%)	83	(0.6%)	
Other/Unknown Pediatric	(0.2%)	8	(0.2%)	1	(0.3%)	22	(0.2%)	30	(0.2%)	
Missouri Total422	(100.0%)	4,532	(100.0%)	370 (	100.0%)	9,119	(100.0%)	13,651	(100.0%)	

\*HIV Cases-Persons with HIV infection who have not developed one of the specific diseases or conditions which would cause them to meet the case definition for AIDS.


\*\*AIDS Cases-Persons with HIV infection who have developed one or more of the specific diseases or conditions which cause them to meet the AIDS case definition.

MSM=men who have sex with men; MSM/IDU=men who have sex with men and inject drugs; IDU=injecting drug users

#### Reported Adult/Adolescent HIV Cases by Exposure Category<sup>1</sup>, Missouri, 1985-2001



### Reported Adult/Adolescent AIDS Cases by Exposure Category<sup>1</sup>, Missouri, 1982-2001



MSM=men who have sex with men; MSM/IDU=men who have sex with men and inject drugs; IDU=injecting drug users; HETSX=heterosexual contact; Oth/Unk=Other/Unknown

# Reported HIV and AIDS Cases and Rates by Area of Residence at Time of Diagnosis, Missouri, 1982–2001

		HIV Cases*				AIDS Cases**				
	F	Reported 2001		Cumulative		Reported 2001		Cumulative		
Geographic Area	Cases	%	Rate***	Cases	%	Cases	%	Rate***	Cases	%
Location										
St. Louis City <sup>†</sup>	135	(32.0%)	40.4	1,326	(29.3%)	140	(37.8%)	41.9	2,582	(28.3%)
St. Louis County <sup>†</sup>	64	(15.2%)	6.4	599	(13.2%)	64	(17.3%)	6.4	1,416	(15.5%)
Kansas City <sup>†</sup>	89	(21.1%)	20.3	1,131	(25.0%)	73	(19.7%)	16.7	2,535	(27.8%)
Outstate <sup>†</sup>	104	(24.6%)	2.8	1,154	(25.5%)	85	(23.0%)	2.3	2,368	(26.0%)
Missouri Correctional Facil	ities †† 30	(7.1%)		322	(7.1%)	8	(2.2%)		218	(2.4%)
Missouri Total	422	(100.0%)	7.7	4,532 (	(100.0%)	370 (	100.0%)	6.8	9,119	(100.0%)


<sup>\*</sup>HIV Cases-Persons with HIV infection who have not developed one of the specific diseases or conditions which would cause them to meet the case definition for AIDS.

<sup>\*\*</sup>AIDS Cases-Persons with HIV infection who have developed one or more of the specific diseases or conditions which cause them to meet the AIDS case definition.

<sup>\*\*\*</sup>Per 100,000 population, based on 1999 population estimates.

<sup>†</sup>Does not include persons living in correctional facilities at the time of diagnosis. These persons are included in the "Missouri Correctional Facilities" category.

 $<sup>\</sup>ensuremath{^{\dagger\dagger}}\xspace$  Includes state, county and local correctional facilities.



### Reported HIV Cases by Race/Ethnicity and Area of Residence at Time of Diagnosis, Missouri, 2001

	Tota	l	White	•	Blaci	<b>(</b>	Hispan	ic
Georgraphic Area	2001 Cases	Rate*	2001 Cases	Rate*	2001 Cases	Rate*	2001 Cases	Rate*
St. Louis City†	135	40.4	39	26.5	90	.51.1	2	34.1
St. Louis County <sup>†</sup>	64	6.4	28	3.5	34	. 20.6	1	7.3
Kansas City†	89	20.3	40	14.3	45	. 34.6	4	21.2
Outstate†	104	2.8	81	2.3	16	. 11.4	3	5.7
Missouri Correctional Facilities <sup>†</sup>	†30		13		17		0	
Missouri Total	422	7.7	201	4.3	202	33.1	10	.10.9


<sup>\*</sup>Per 100,000 population, based on 1999 population estimates.

Currently Living HIV-Diagnosed Persons (HIV and AIDS Cases) Who Were Residents of Missouri at the Time of Diagnosis, and


Who Were Reported Through 2001, by Gender and Race/Ethnicity


	Living HIV-
	Diagnosed Persons %
Gender	
Male	7.336 85.1%
Female	
Race/Ethnicity	
White	4.845 56.2%
Black	*
Hispanic	
Asian/Pacific Islander	
American Indian	
Unknown	
Race/Ethnicity and Gend	er
White Male	4.361 50.6%
BlackMale	
Hispanic Male	
Asian/Pacific Islander Male	20 0.2%
American Indian Male	27 0.3%
Unknown Male	17 0.2%
WhiteFemale	
Black Female	
HispanicFemale	23 0.3%
Asian/Pacific Islander Female	
American Indian Female	
Unknown Female	4 0.0%
Total Living HIV-Diagnosed I	Persons8.616100.0%

Currently Living HIV-Diagnosed Persons (HIV and AIDS Cases), Reported Through 2001, by Missouri County of Residence † at Time of Diagnosis




<sup>†</sup>Does not include persons living in correctional facilities at the time of diagnosis. These persons are included in the "Missouri Correctional Facilities" category. ††Includes state, county and local correctional facilities.







### Reported Gonorrhea Cases and Rates by Geographic Area, Missouri, 2001

Cases	s %	Rate*
Missouri		
Whites 1,098	12.6%	23.4
Blacks 6,562	75.2%	1,074.0
Other/Unknown 1,063	12.2%	
Total Cases 8,723	100.0%	159.5
St. Louis City		
Whites 132	4.1%	89.7
Blacks 2,707	85.0%	1,535.8
Other/Unknown346	10.9%	
Total Cases 3,185	100.0%	953.7
St. Louis County		
Whites 120	6.5%	15.1
Blacks	80.3%	900.6
Other/Unknown243	13.2%	
Total Cases 1,847	100.0%	185.4
Kansas City		
Whites 181	9.0%	64.8
Blacks 1,672	82.6%	
Other/Unknown171	8.4%	1,204.3
Total Cases 2,024		462.3
10441 04505 111111 2,021	200.070	.02.0
Outstate		
Whites 665	39.8%	19.2
Blacks 699	41.9%	500.8
Other/Unknown303	18.2%	
Total Cases 1,667	100.0%	45.1
*Per 100,000 population		










### Reported P&S Syphilis Cases and Rates by Geographic Area, Missouri, 2001


Cases	%	Rate*
Missouri		
Whites5	19.2%	0.1
Blacks19	73.1%	3.1
Other/Unknown2	7.7%	
Total Cases26	100.0%	0.5
St. Louis City		
Whites1	6.7%	0.7
Blacks 14	93.3%	7.9
Other/Unknown0	0.0%	
Total Cases15	100.0%	4.5
St. Louis County		
Whites0	0.0%	0.0
Blacks 0	0.0%	0.0
Other/Unknown1	100.0%	
Total Cases1	100.0%	0.1
Kansas City		
Whites2	28.6%	0.7
Blacks 3	57.1%	2.3
Other/Unknown0	14.3%	
Total Cases5	100.0%	1.6
Outstate		
Whites2	40.0%	0.1
Blacks2	40.0%	1.4
Other/Unknown1	10.0%	
Total Cases5	100.0%	0.1
*Per 100,000 population		





Congenital Syphilis: 5 Congenital Syphilis cases were reported in 2001
5 (100.0%) African American
4 (80.0%) St. Louis City
1 (10.0%) St. Louis County





### Reported Chlamydia Cases and Rates by Geographic Area, Missouri, 2001

	Cases	%	Rate*
Missouri			
Whites	4,563	32.7%	97.3
Blacks		45.9%	1,047.5
Other/Unknown .	2,986	21.4%	
Total Cases	.13,949	100.0%	255.1
St. Louis City			
Whites	162	5.1%	110.1
Blacks		73.0%	1,323.6
Other/Unknown .		21.9%	
Total Cases		100.0%	956.7
Total Cases	0,170	100.070	,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,
St. Louis Count			
Whites	338	13.2%	42.4
Blacks		64.9%	1,008.6
Other/Unknown .	560	21.9%	
Total Cases	2,560	100.0%	257.0
Kansas City			
Whites	458	19.3%	164.0
Blacks	1,453	61.4%	1,116.1
Other/Unknown .	456	19.3%	
Total Cases	2,367	100.0%	540.7
Outstate			
Whites	3.607	61.8%	104.1
Blacks		16.4%	683.3
Other/Unknown .		21.8%	
Total Cases		100.0%	157.5
*Per 100,000 population	,==-		

